

BPS ansatzes as electric form-factors.

L. D. Lantsman.

18109, Rostock, Germany; Mecklenburger Allee, 7

llantsman@freenet.de

Tel. (049)-0381-7990724.

February 6, 2009

Abstract

We argue that BPS ansatzes, entering manifestly vacuum BPS monopole solutions to equations of motion in the (Minkowskian) non-Abelian Higgs model play the role of some electric form-factors and that this implies (soft) violating the CP-invariance of the mentioned model, similar to taking place in the Euclidian Yang-Mills (YM) theory with instantons, generating the θ -term in the appropriate effective Hamiltonian.

PACS: 14.80.Bn, 14.80.Hv.

Keywords: Non-Abelian Theory, BPS Monopole, Minkowski Space, Instanton.

Contents

1	Introduction.	3
2	BPS ansatzes as electric form-factors.	7
3	Higgs BPS ansatzes and CP violating.	10
4	Discussion.	14

1 Introduction.

The (without of quarks) non-Abelian YM theory involving vacuum BPS monopole solutions in their Higgs and gauge sectors (as a result of the spontaneous breakdown the initial $SU(2)$ gauge symmetry group to its $U(1)$ subgroup) occupy a special position among another such theories with monopoles. This is associated with manifest superfluid properties of the former model.

To elucidate this our assertion, let us at first write down explicitly the action functional for the (Minkowskian) Yang-Mills-Higgs (YMH) model. It can be represented as [1, 2, 3, 4, 5]

$$S = -\frac{1}{4g^2} \int d^4x F_{\mu\nu}^b F_b^{\mu\nu} + \frac{1}{2} \int d^4x (D_\mu \phi, D^\mu \phi) - \frac{\lambda}{4} \int d^4x \left[(\phi^b)^2 - \frac{m^2}{\lambda} \right]^2, \quad (1.1)$$

with

$$D_\mu \phi = \partial^\mu \phi + g[A^\mu, \phi]$$

being the covariant derivative and g being the YM coupling constant.

The action functional (1.1) results the equations of motion [1]

$$(D_\nu F^{\mu\nu})_a = -g\epsilon_{abc}\phi^b (D_\mu \phi)^c, \quad (1.2)$$

$$(D^\mu D_\mu \phi)_a = -\lambda \phi_a (\vec{\phi} \cdot \vec{\phi} - a^2); \quad a^2 = m^2/\lambda. \quad (1.3)$$

It turns out that going over to the limit

$$\lambda \rightarrow 0, \quad m \rightarrow 0: \quad \frac{1}{\epsilon} \equiv \frac{gm}{\sqrt{\lambda}} = ga \neq 0; \quad (1.4)$$

in Eq. (1.1) just induces the (topologically degenerated) vacuum BPS monopole solutions in the Higgs and YM sectors of the model (1.1).

Historically, the idea to go over to the limit (1.4) in the YMH model is originated from the works [6], and from this time it was referred to as the *Bogomol'nyi-Prasad-Sommerfeld* (BPS) limit. Later, in the papers [3, 4] the BPS limit was rearranged to the look (1.4), implicating the YM coupling constant g .

It is remarkable that the ratio $a = m/\sqrt{\lambda}$ (having the mass dimension) can take arbitrary values in the limit (1.4) and that the variable $\epsilon \rightarrow 0$ of the length dimension is introduced therein. We shall make sure soon that ϵ plays the role of the size parameter characterized the core of a BPS monopole.

Vacuum BPS monopole solutions can be derived in the limit (1.4) at evaluating the lowest bound of the energy for the given YMH configuration (often referred to as the *Bogomolnyi bound* in the modern literature) [3, 4]:

$$E_{\min} = 4\pi \mathbf{m} \frac{a}{g} \quad (1.5)$$

(where $\mathbf{m}$ denotes the magnetic charge).

As a result, one arrives at the so-called *Bogomolnyi equation* [3, 4, 7]

$$\mathbf{B}(\Phi) = \pm D\Phi \quad (1.6)$$

relating the vacuum “magnetic” field $\mathbf{B}$ to the vacuum Higgs configuration in the shape of a BPS monopole. The presence of two opposite signs in the Bogomolnyi equation (1.6) corresponds to two opposite signs of magnetic charges in nature.

The explicit way deriving the Bogomolnyi equation (1.6) and evaluating the Bogomolnyi bound (1.5) was stated, for instance, in the monograph [8] (see *ibid* §Φ11). In particular, in the monograph [8] vacuum BPS monopole solutions to the Bogomolnyi equation (1.6), arising in the Higgs and gauge sectors of the YMH model [6], were written down. In the series of papers [3, 4] these solutions were reproduced with

the only modification that the effective Higgs mass $a = m/\sqrt{\lambda}$ (utilized in Ref. [8]) was replaced with the parameter ϵ^{-1} :

$$\Phi_{(0)}^a(t, \mathbf{x}) = \frac{x^a}{gr} f_0^{BPS}(r), \quad f_0^{BPS}(r) = \left[\frac{1}{\epsilon \tanh(r/\epsilon)} - \frac{1}{r} \right], \quad (1.7)$$

$$A_{i(0)}^a(t, \mathbf{x}) \equiv \Phi_i^{aBPS}(\mathbf{x}) = \epsilon_{iak} \frac{x^k}{gr^2} f_1^{BPS}(r), \quad f_1^{BPS}(r) = \left[1 - \frac{r}{\epsilon \sinh(r/\epsilon)} \right]. \quad (1.8)$$

Indeed, the BPS monopoles (1.7), (1.8) are topologically trivial fields ¹.

Now let us discuss the behaviour of the *BPS ansatzes* $f_0^{BPS}(r)$ and $f_1^{BPS}(r)$ at the origin of coordinates and at the spatial infinity. Direct checking shows that [13]

$$f_0^{BPS}(0) = 0, \quad f_0^{BPS}(\infty) = 1; \quad (1.9)$$

$$f_1^{BPS}(0) = 0, \quad f_1^{BPS}(\infty) = 1. \quad (1.10)$$

Then the YM BPS monopoles (1.8) (with their Gribov copies $\Phi_i^{a(n)}$) display an alike good behaviour (disappearing) at the origin of coordinates and at the spatial infinity ($r \rightarrow \infty$). The other thing the behaviour of Higgs vacuum BPS monopole modes $\Phi_{(n)a}$. These diverge at the origin of coordinates, as it follows from (1.7) and (1.9).

The interesting and important feature of YM BPS monopoles (1.8) is that they merge (because of Eq. (1.10)) with Wu-Yang monopoles Φ_i^{Wa} [14] ².

¹The topological degeneration of vacuum BPS monopole data (1.7), (1.8) can be carry out by means of “large” gauge transformations (in the terminology [9]) proposed in the work [10]:

$$\Phi_i^{a(n)} := v^{(n)}(\mathbf{x}) [\Phi_i^{a(0)} + \partial_i] v^{(n)}(\mathbf{x})^{-1}, \quad v^{(n)}(\mathbf{x}) = \exp[n\hat{\Phi}_0(\mathbf{x})];$$

$$\Phi_{(n)a} = v^{(n)}(\mathbf{x}) \Phi_{(0)a} v^{(n)}(\mathbf{x})^{-1}; \quad n \in \mathbf{Z}$$

(latter Eq. was derived in the paper [11]).

Here

$$\hat{\Phi}_0(r) = -i\pi \frac{\tau^a x_a}{r} f_{01}^{BPS}(r); \quad f_{01}^{BPS}(r) = \left[\frac{1}{\tanh(r/\epsilon)} - \frac{\epsilon}{r} \right] = f_1^{BPS}(r)/\epsilon;$$

with τ^a ($a = 1, 2, 3$) being the Pauli matrices. The exponential multipliers $v^{(n)}(\mathbf{x})$ were referred to as *Gribov topological multipliers* in Ref. [10] while the value $\hat{\Phi}_0(r)$ as the *Gribov phase*.

It can be argued that specified in this way topologically degenerated YM vacuum BPS monopole data $\Phi_i^{a(n)}$ satisfy the Coulomb gauge $D^i \Phi_i^{a(n)} = 0$. And moreover, YM vacuum BPS monopole data $\Phi_i^{a(n)}$ also turn out to be gauge invariant, i.e. *physical*, functionals of YM fields. In particular, this is correctly for topologically trivial YM BPS monopoles (1.8) [3, 4], and this indicates transparently the purely physical nature of these monopoles. Also topologically trivial Higgs BPS monopole modes (1.7) and their Gribov topological copies $\Phi^{(n)a}$ [11] prove to be manifestly gauge invariant, side by side with YM BPS monopoles.

Mention that the topologically degenerated YM vacuum BPS monopoles $\Phi_i^{a(n)}$ are patterns of *topological Dirac variables*, gauge invariant and transverse (in the sense satisfying the Lorentz covariant Coulomb gauge $D^i \Phi_i^{a(n)} = 0$).

The important point here that topological Dirac variables $\Phi^{(n)a}$ are got indeed as solutions to the YM Gauss law constraint [10]

$$\frac{\delta W}{\delta A_0^a} = 0 \iff [D^2(A)]^{ac} A_{0c} = D_i^{ac}(A) \partial_0 A_c^i.$$

As to Higgs BPS monopole modes $\Phi^{(n)a}$, the appropriate current $\rho^H \sim ig\Phi D\Phi$ decouples from the YM Gauss law constraint in the first order of the perturbation theory by the YM coupling constant g .

The detailed analysis of topological Dirac variables (including the answer why Higgs BPS monopole modes $\Phi^{(n)a}$ disappear from the YM Gauss law constraint in the first order of the perturbation theory) was performed in the works [3, 4, 10, 12, 13], and we recommend these to our readers for studying the matter.

²Remember that Wu-Yang monopoles Φ_i^{Wa} are solutions to the classical equation of motion [3, 4, 10]

$$D_k^{ab}(\Phi_i^W) F_a^{bk}(\Phi_i^W) = 0 \implies \frac{d^2 f}{dr^2} + \frac{f(f^2 - 1)}{r^2} = 0$$

in the “pure” YM theory (with absent Higgs and fermionic modes) corresponding to the exact $SU(2)$ gauge group.

The Bogomolnyi equation (1.6) can be treated as a *potentiality condition* for the BPS monopole vacuum. A brief argumentation in favour of this statement was advanced in the recent paper [5].

Really, mathematically, any potentiality condition may be written down as

$$\text{rot grad } \Phi = 0 \quad (1.11)$$

for a scalar field Φ . Thus any potential field may be represented as $\text{grad } \Phi$.

In the Minkowskian YMH theory involving BPS monopole solutions there exists always such scalar fields. There are just Higgs vacuum BPS monopole modes (1.7) (with their Gribov topological copies $\Phi_{(n)a}$ [11]).

Then it is easy to guess that the Bogomol'nyi equation (1.6), having the look (1.11), can be treated as the potentiality condition for the Minkowskian YMH vacuum involving vacuum BPS monopole solutions. It is so due to the Bianchi identity $DB = 0$ ³.

Indeed, there can be drawn a highly transparent parallel between the Minkowskian YMH vacuum involving vacuum BPS monopole solutions and a liquid helium II specimen described in the Bogolubov-Landau model [17].

In the latter case, the potential motion is proper to the superfluid component in this liquid helium specimen.

The superfluid motion in a liquid helium II is the motion without a friction between the superfluid component and the walls of the vessel where a liquid helium specimen is contained.

Thus the viscosity of the superfluid component in a helium II is equal to zero, and vortices (involving $\text{rot } \mathbf{v} \neq 0$) are absent in the superfluid component of a helium.

As L. D. Landau showed [17], at velocities of the liquid exceeding a *critical velocity* $v_0 = \min(\epsilon/p)$ for the ratio of the energy ϵ and momentum p for quantum excitations spectrum in the liquid helium II, the dissipation of the liquid helium energy occurs via arising excitation quanta with momenta $\mathbf{p}$ directed antiparallel to the velocity vector $\mathbf{v}$. Such dissipation of the liquid helium energy becomes advantageous [18] just at

$$\epsilon + \mathbf{p} \cdot \mathbf{v} < 0 \implies \epsilon - p v < 0.$$

From the above reasoning concerning properties of potential motions, it becomes obvious that the vector $\mathbf{v}_0$ of the critical velocity for the superfluid potential motion possesses the zero curl: $\text{rot } \mathbf{v}_0 = 0$.

In this case, according to (1.11), the critical velocity $\mathbf{v}_0$ of the superfluid potential motion in a liquid helium specimen may be represented [19] as

$$\mathbf{v}_0 = \frac{\hbar}{m} \nabla \Phi(t, \mathbf{r}), \quad (1.12)$$

where m is the mass of a helium atom and $\Phi(t, \mathbf{r})$ is the phase of the complex-value helium Bose condensate wave function $\Xi(t, \mathbf{r}) \in C$.

Thus the similar look for the vacuum "magnetic" field $\mathbf{B}$ in the Minkowskian Higgs model involving BPS monopole solutions, generating by the Bogomol'nyi equation (1.6), and for the critical velocity $\mathbf{v}_0$

One can distinguish three solutions to this equation:

$$f_1^{PT} = 0, \quad f_1^W = \pm 1 \quad (r \neq 0).$$

The first, trivial, solution $f_1^{PT} = 0$ corresponds to the naive unstable perturbation theory, involving the asymptotic freedom formula [15, 16]. They are just the Wu-Yang monopoles [14] with topological charges ± 1 , respectively.

³This becomes more transparent upon representing the Bogomol'nyi equation (1.6) in the tensor shape [8]

$$\frac{1}{2g} \epsilon^{ijk} F_{jk}^a = \nabla^i \Phi^a.$$

Then due to the Bianchi identity

$$\epsilon^{ijk} \nabla_i F_{jk}^b = 0,$$

the Bogomol'nyi equation (1.6) results

$$D^2 \Phi \sim \text{rot } \mathbf{B} = 0$$

(at neglecting the items in DB directly proportional to g and g^2).

of the superfluid motion in a liquid helium II, given by Eq. (1.12), testifies in favour of the potential motions occurring therein.

In this case, drawing a highly transparent parallel between the Minkowskian YMH vacuum involving BPS monopole solutions and a liquid helium II specimen described in the Bogolubov-Landau model [17], we can also conclude about manifest superfluid properties of the Minkowskian YMH vacuum involving BPS monopoles.

As in the Bogolubov-Landau model [17] of liquid helium II, the ground cause of the superfluid properties of the Minkowskian YMH vacuum with BPS monopoles roots in long-range correlations of local excitations [20].

While in the Bogolubov-Landau model [17] of liquid helium II this comes to repulsion forces between helium atoms as the cause of superfluidity effects, in the Minkowskian YMH vacuum involving BPS monopole solutions, the cause of the superfluidity taking place is in the strong YMH coupling g (entering effectively the appropriate action functional (1.1)).

The principal thing in alike superfluid effects occurring in a liquid helium II specimen as well as in the Minkowskian YMH vacuum involving BPS monopoles is that these both physical systems are *non-ideal gases*.

In ideal gases no superfluidity phenomena are possible.

There can be demonstrated [21] that in ideal gases a deal of particles is accumulated on the zero energy quantum level at temperatures $T < T_0$; herewith the temperature T_0

$$kT_0 = \frac{1}{(2.61)^{2/3}} \frac{h^2}{2\pi m} \left(\frac{N}{V}\right)^{2/3} \quad (1.13)$$

(with k and h being, respectively, the Boltzmann and Planck constants; N being the complete number of particles; V being the volume occupied by the ideal Bose gas; m being the mass of a particle) is called *the condensation temperature*, while the above deal of particles is called *the Bose condensate*.

The just described superfluidity is absent in another Minkowskian Higgs models with monopoles: for instance, in the 't Hooft-Polyakov model [22, 23]. This can be argued, repeating the arguments [5, 24], by disappearing the covariant derivative $D^i \phi_a$ of a Higgs 't Hooft-Polyakov monopole mode ϕ_a at the spatial infinity.

In this case, asymptotically (at $r \rightarrow \infty$),

$$\mathbf{B}_i^a D^i \phi_a = \partial_i (\mathbf{B}_a^i \phi_a) = 0, \quad (1.14)$$

because of the Bianchi identity $DB = 0$ and the remark that $\mathbf{B}_a^i \phi_a$ is a $U(1) \subset SU(2)$ scalar; thus one can replace the covariant derivative D with the partial one, ∂ , for $\mathbf{B}_a^i \phi_a$.

In turn, the complete energy of the YMH configuration may be represented as [8, 24]

$$E_{\text{compl}} = \int d^3x \left[\frac{1}{2} (D\phi_a \pm \mathbf{B}_a)^2 + \frac{\lambda}{4} ((\phi^a)^2 - a^2) \right] + \frac{4\pi}{g^2} M_W. \quad (1.15)$$

The last item in Eq. (1.15) involves the mass M_W of the W -boson.

Such look of E_{compl} originates from the paper [22] devoted to the 't Hooft-Polyakov model.

The connection between the energy integral E_{compl} and the general action functional (1.1) [3, 4] of the Minkowskian Higgs model is given by the identity [24]

$$(D\phi_a)^2 + \mathbf{B}_a^2 = (D\phi_a \pm \mathbf{B}_a)^2 \mp 2\mathbf{B}_a D\phi_a. \quad (1.16)$$

Herewith the last item on the right-hand side of (1.16) vanishes at the spatial infinity, as we have noted above. Just from Eq. (1.15) one can read the Bogomol'nyi equation in the shape (1.6).

In the 't Hooft-Polyakov model [22, 23] the Bogomol'nyi equation (1.6) determines the Bogomol'nyi bound [24]

$$M_{\text{mon}} = \frac{4\pi}{g^2} M_W \quad (1.17)$$

for the complete energy E_{compl} , (1.15), of the YMH configuration at going over to the BPS limit (1.4) [6].

Then the asymptotic $D_i\phi^a \rightarrow 0$ as $r \rightarrow \infty$ for 't Hooft-Polyakov monopoles [22, 23] forces to vanish identically the first item under the integral sign in E_{compl} ($|\mathbf{B}| = 0$).

In the light of the said above it becomes obvious that the vacuum "magnetic" field $\mathbf{B}$, playing the role of the (critical) velocity for the superfluid motion in the Minkowskian non-Abelian vacuum with BPS monopoles, actually approaches zero in the 't Hooft-Polyakov model [22, 23], involving the $D_i\phi^a \rightarrow 0$ as $r \rightarrow \infty$ asymptotic for Higgs monopoles.

The principal goal of the present note is to show that BPS ansatzes $f_1^{BPS}(r)$, $f_0^{BPS}(r)$, one encounter in the Higgs BPS monopole model, can serve as electric form-factors therein. Unlike this, electric form-factors become trivial in the 't Hooft-Polyakov theory [22, 23]. Grounding this fact will be the topic of Section 2.

In Section 3 we show that presence of BPS ansatzes in the considered model implies violating the CP invariance. It is the effect similar to that taking place in the instanton models [1, 2, 8, 25], generating the θ -items in the appropriate effective Lagrangians. This effect violating the CP invariance by the θ -dependence of the instanton models was analyzed in the paper [26], and we reconstruct partially the arguments [26] in Section 3.

2 BPS ansatzes as electric form-factors.

The starting point of our discussion will be the well known Dirac quantization condition [27] for the electric and magnetic charges presented in a closed system of quantum fields.

In a simple case when a quantum object is isolated from another, the Dirac quantization condition [27] acquires the look [1, 2, 8]

$$\frac{q\mathbf{m}}{4\pi} = \frac{1}{2}n; \quad n \in \mathbf{Z}, \quad (2.1)$$

where q and $\mathbf{m}$ are, respectively, the electric and magnetic charges of the considered object (in the system of units in which $\hbar = c = 1$)⁴. Each such quantum object possessing the electric and magnetic charges simultaneously is referred to as a *dyon* in modern physical literature⁵.

When a system of quantum fields consists of two dyons, Eq. (2.1) can be generalized to Eq. [24, 26]

$$q_1\mathbf{m}_2 \pm q_2\mathbf{m}_1 = 2\pi n; \quad n \in \mathbf{Z}; \quad (2.2)$$

⁴In some sources (for instance, [2, 8]) Eq. (2.1) is given in the slightly modified air

$$q\mathbf{m} = \frac{1}{2}n.$$

Going over from Eq. (2.1) to the latter Eq. can be achieved [24] at setting $\mathbf{m} = 4\pi/q$. The origin of this in the Laplace equation [2]

$$\nabla \cdot \mathbf{B} = 4\pi\mathbf{m}\delta^3(r)$$

for the point magnetic charge $\mathbf{m}$ creating the radial magnetic field $\mathbf{B}$, resulting the total magnetic flux

$$\Phi = 4\pi r^2 B = 4\pi\mathbf{m}$$

through a sphere with its centre in the origin of coordinates.

Just this provides (see §10.3 in [2]) the change

$$\Delta\alpha|_\pi = \frac{q}{\hbar c} \oint \mathbf{A} \cdot d\mathbf{l}|_\pi = \frac{q}{\hbar c} \int \text{rot}\mathbf{A} \cdot d\mathbf{S}|_\pi = \frac{q}{\hbar c} \int \mathbf{B} \cdot d\mathbf{S}|_\pi = \frac{q}{\hbar c} \Phi(r, \theta)|_{\theta=\pi} = \frac{q}{\hbar c} 4\pi\mathbf{m} = 2\pi n$$

of the dyon's wave function ($\psi \equiv |\psi|e^{i\alpha} = |\psi|\exp[(-iq/\hbar c)\mathbf{A} \cdot \mathbf{r}]$) phase α at $\theta = \pi$ (the flux $\Phi(r, \pi)$ is just the maximal possible flux, spreaded to the whole sphere).

⁵Formally, a particle possessing the zero electric and magnetic charges also relates to the class of dyons. And moreover, if $\mathbf{m} = 0$, Eq. (2.1) is satisfied at arbitrary values of the electric charge q .

Eq. (2.2) was derived for the first time by Zwanziger and Schwinger [28]. The reasoning for deriving this Eq. is [26] the classical formula for the angular momentum of an electromagnetic field. The angular momentum in an electromagnetic background of a two-particle system can be calculated easily. It has the magnitude

$$(e_1 \mathbf{m}_2 - e_2 \mathbf{m}_1)/4\pi c,$$

that takes integer or half-integer values, as it is expected in quantum mechanics, only if

$$(e_1 \mathbf{m}_2 - e_2 \mathbf{m}_1)/\hbar c = 2\pi n$$

(with setting $\hbar = c = 1$). Going over to the sign + in (2.2) from the – one is reduced simply to replacing $\mathbf{m} \leftrightarrow -\mathbf{m}$.

In Ref. [24] it was given the in definite sense generalization of Eq. (2.2) to the case of an arbitrary gauge group $SU(N)$:

$$\sum_{i=1}^{N-1} e_i \mathbf{m}_i = 2\pi n.$$

Herewith it is easy to see that Eq. (2.2) (with the + sign) is the particular case of the latter relation for the gauge group $SU(2)$.

Let us now calculate, following [1], the total momentum

$$\mathbf{J} = \mathbf{L} + \mathbf{T} \quad (2.3)$$

of a particle in a magnetic monopole background. It involves its spatial angular momentum $\mathbf{L}$ (including its "ordinary" spin) and the generator $\mathbf{T}$ of the internal (for instance, the gauge $U(1)$) symmetry.

On the other hand, $\mathbf{L} = \mathbf{r} \times \mathbf{p}$, with $\mathbf{p}$ being the canonical momentum

$$p_i = mv_i + g(\mathbf{A}_i \cdot \mathbf{T}) \quad (2.4)$$

for a (vacuum) "gauge" monopole solution involving the mass m .

In the particular case of 't Hooft-Polyakov monopole solutions [22, 23], when YM potentials have the look [2]

$$A_i^a = \epsilon_{iab} \frac{r^b}{gr^2}, \quad (2.5)$$

Eq. (2.4) can be rewritten as

$$\mathbf{p} = m\dot{\mathbf{r}} + \frac{1}{r} \mathbf{n} \times \mathbf{T}. \quad (2.6)$$

Then

$$\begin{aligned} \mathbf{J} &= \mathbf{r} \times \mathbf{p} + \mathbf{T} = \mathbf{r} \times m\dot{\mathbf{r}} + \mathbf{n} \times \mathbf{n} \times \mathbf{T} + \mathbf{T} = \\ &= \mathbf{r} \times m\dot{\mathbf{r}} + (\mathbf{n} \cdot \mathbf{T})\mathbf{n}. \end{aligned} \quad (2.7)$$

Now we must recall that in the 't Hooft-Polyakov model [22, 23] the radial "magnetic" field $\mathbf{B}$ is given by Eq. [2]⁶

$$F^{ij} = -\frac{1}{gr^3} \epsilon^{ijk} r_k; \quad \mathbf{B}_k \equiv -F^{ij} \epsilon_{ijk}. \quad (2.8)$$

From the general reasoning [1] about magnetic monopoles, the equation of motion for an electric charged particle in its field is read as

$$m\ddot{\mathbf{r}} = e\dot{\mathbf{r}} \times \mathbf{B}. \quad (2.9)$$

It is just the Lorenz force acting onto this particle in the magnetic monopole background.

⁶Note that Eqs. (2.5) and (2.8) correspond to the "standard" normalization YMH Lagrangian density [2] involving the coefficient $-1/4$ in front of $F_{\mu\nu}^2$.

In this case the rate of change of the particle angular momentum $\mathbf{L}$ is

$$\frac{d}{dt}(\mathbf{r} \times m\dot{\mathbf{r}}) = \mathbf{r} \times m\ddot{\mathbf{r}} = \frac{2\pi n}{\nu} \mathbf{r} \times (\dot{\mathbf{r}} \times \mathbf{r}) = \frac{d}{dt} \left(\frac{2\pi n}{\nu} \mathbf{n} \right), \quad (2.10)$$

where [8] ν is the (minimal) positive number for which the condition $\exp(\nu h) = 1$ (with $h \equiv h(\Phi) \equiv \Phi/a$ being the generator of the residual $U(1)$ gauge group in the quested YMH model) is satisfied ⁷.

Then, issuing from the above discussed Dirac quantization condition

$$q\mathbf{m} = \frac{1}{2}n \quad (2.11)$$

and the normalizations [8]

$$q = \frac{2\pi n}{\nu} g, \quad (2.12)$$

$$\mathbf{m} = \frac{\nu}{4\pi g} \zeta; \quad \zeta \in \mathbf{Z}; \quad (2.13)$$

for the electric and magnetic charges, respectively (thus the Dirac quantization condition (2.11) is satisfied automatically), we just arrive to Eq. (2.10) (with q given in (2.12) and appropriate cancelling the YM coupling constant g , which enters the relation for $\mathbf{B}$).

The said suggests the formal possibility to introduct the total momentum (2.3) [1] of an electric charged particle in the magnetic monopole background in such a wise that it is conserved:

$$\mathbf{J} = \mathbf{r} \times m\dot{\mathbf{r}} - \frac{2\pi n}{\nu} \mathbf{n} = \mathbf{r} \times m\dot{\mathbf{r}} - (q/g)\mathbf{n}. \quad (2.14)$$

Comparing then the expressions (2.7) and (2.14) for the total momentum $\mathbf{J}$ of an electric charged particle in the magnetic monopole background got in the 't Hooft-Polyakov model [22, 23], one can conclude that

$$(e/g) = -\mathbf{n} \cdot \mathbf{T} \quad (2.15)$$

if $q = e$ (e is the elementary charge).

At the particular choice $\mathbf{n}$ to be the z -direction, $\mathbf{n} \cdot \mathbf{T} = T_3$. On the other hand, the value $2\pi/\nu$ can be normalized as $2\pi/\nu = 1$ (at considering [8] the $U(1)$ group space as the circle S^1 of the unit radius).

Because of (2.12), we can conclude that the isospin operator $\mathbf{T}$ (T_3) is topologically degenerated. Geometrically, such topological degeneration of the isospin operator $\mathbf{T}$ means extracting ("large" and "small") gauge orbits in the $U(1)$ group space.

Specifying [1] the electric charge operator $Q_{U(1)} = eT_3$, we see additionally that $Q_{U(1)}$ takes integers multipliers of e (at setting $2\pi/\nu = 1$) in the presence of 't Hooft-Polyakov monopole modes ⁸.

The presence of YM BPS ansatz (1.8) in the YMH model with (vacuum) BPS monopole solutions changes the computations [1] regarding the isospin operator $\mathbf{T}$ and the total momentum $\mathbf{J}$.

So instead of (2.7), then it should be written down

$$\mathbf{J} = \mathbf{r} \times m\dot{\mathbf{r}} + f_1^{BPS}(\mathbf{n} \cdot \mathbf{T})\mathbf{n} + \mathbf{T}(1 - f_1^{BPS}). \quad (2.16)$$

⁷Defining ν in this way, one can argue [8] that

$$\mathbf{m}(\Phi, A) = C \zeta(\Phi, A), \quad \zeta(\Phi, A) \in \mathbf{Z}$$

for the given monopole YMH configuration (Φ, A) with $C = \nu/4\pi$.

⁸This effect was noted, for example, in Ref. [24] with that important correction that besides $Q_{U(1)} = eT_3 = ne$ charged states, $(n + 1/2)e$ states are also possible, with $q = e/2$ being the minimal charge corresponding to $\mathbf{T} = 1/2$.

For trivial topologies $n = 0$, the minimal charge $q = e/2$ corresponds to a fermionic field ψ in the

$$I = \frac{1}{2} : \quad \psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}$$

representation of $SU(2)$ (if an YMH model is in question).

The third item appearing in (2.16) corresponds to the expression

$$\mathbf{p} = m\dot{\mathbf{r}} + \left(\frac{1}{r}f_1^{BPS}\right)\mathbf{n} \times \mathbf{T}$$

for the momentum $\mathbf{p}$ of a particle in the YM BPS monopole background.

Indeed, the presence of the YM BPS ansatz (1.8) complicates to a considerable extent the computations comparing to those (2.10)- (2.14) [1] in the 't Hooft-Polyakov monopole model [22, 23]. It is associated, for instance, with the more complicated expression for $\mathbf{B}$ (see e.g. [7]) in the BPS monopole theory.

But, in spite of these difficulties, one can conclude, issuing from (2.16), that f_1^{BPS} really plays the role of an electric form-factor in the BPS monopole YMH model. It is because the second item in (2.16) can be represented as

$$f_1^{BPS}T_3\mathbf{n},$$

that implies, as it is easy to understand, the replacement $e \leftrightarrow f_1^{BPS}e$ in Eq. (2.16). And this is just equivalent to the role of the YM BPS ansatz f_1^{BPS} , (1.8), as an electric form-factor screening the (elementary) charge e .

The similar role of an electric form-factor is played also by the Higgs BPS ansatz f_0^{BPS} , (1.7). The physical consequence of this is screening effect for electric charges in the Higgs phase [24] of the YMH model, additional to that rendering in the Higgs phase by the in average electrically neutral Higgs Bose condensate.

3 Higgs BPS ansatzes and CP violating.

In this section, repeating the arguments [26], we shall attempt to demonstrate that the presence of YM BPS ansatz f_1^{BPS} , (1.8), in the (Minkowskian) YMH BPS monopole theory violates manifestly the CP invariance of that theory.

In the previous section we have discussed the Dirac-Zwanziger-Schwinger quantization condition (2.2) [26]. It turns out that this condition says something important about the difference between electric charges of two magnetic monopoles.

Given, for example, two monopoles of minimum allowed charge $2\pi/e$ and of electric charges q and q' , one finds

$$e_1\mathbf{m}_2 - e_2\mathbf{m}_1 = 2\pi(q - q')/e, \quad (3.1)$$

so that the Dirac-Zwanziger-Schwinger quantization condition (2.2) gives

$$q - q' = ne. \quad (3.2)$$

Thus the difference $q - q'$ must be an integer multiple of e . But as it was noted in [28], there is no restriction onto q and q' separately.

If, however, the Dirac-Zwanziger-Schwinger quantization condition (2.2) is supplemented by the CP conservation, the allowed values of the electric charge of an magnetic monopole are also quantized.

In fact, although the electric charge is odd under CP, the magnetic charge is even (this is because electric and magnetic fields are transformed oppositely under parity). Applied to a monopole of the charges $(q, 2\pi/e)$, a CP transformation gives the monopole of the charges $(-q, 2\pi/e)$. For these two particles

$$e_1\mathbf{m}_2 - e_2\mathbf{m}_1 = 4\pi q/e \quad (3.3)$$

and is a multiple of 2π only if

$$q = ne \quad \text{or} \quad q = (n + 1/2)e. \quad (3.4)$$

Thus at assuming the CP conservation, monopoles can have integer or half-integer electric charges (as multiples of e). And moreover, if monopoles of integer charges exist, monopoles of half-integer charges do not exist and vice-versa.

Appart from the CP conservation, there are no reasoning for satisfying the claim (3.4). In nature the CP invariance is violated, but weakly. One can thus suspect that monopoles possess almost (half)integer electric charges. The deviation of monopoles from such charges would be proportional to the strenght of CP violating [26].

The one source CP violating is the instanton YM model [25], resulting [1] the effective Lagrangian

$$\mathcal{L}_{\text{eff}} \equiv \mathcal{L} + \Delta\mathcal{L} = \mathcal{L} + \frac{g^2\theta}{16\pi^2} \text{tr} (F_{\mu\nu}^a \tilde{F}^{\mu\nu}); \quad \tilde{F}_{\mu\nu} = \frac{1}{2}\epsilon^{\mu\nu\alpha\beta} F_{\alpha\beta}; \quad (3.5)$$

involving the quasimomentum $\theta \in [-\pi, \pi]$ ⁹.

⁹Indeed, as it was argued in the papers [11, 20, 29] (see also [30]), θ is a complex parameter.

This can be seen at performing the quantization procedure for the instanton YM model [25]. The latter one is reduced to solving the system of equations [11, 20, 29, 30]

$$\begin{aligned} \hat{H}(i\delta/\delta A, A)\Psi_\epsilon[A] &= \epsilon\Psi_\epsilon[A], \\ \nabla_i^{ab}(A)\left(\frac{\delta}{i\delta A_{ib}}\Psi_\epsilon[A]\right) &= 0; \quad \nabla_i^{ab}(A) = \delta^{ab}\partial_i - g\epsilon^{abc}A_{ic}; \\ T_1\Psi_\epsilon[A] &= e^{i1\cdot\theta}\Psi_\epsilon[A]. \end{aligned}$$

for the wave function $\Psi_\epsilon[A]$, the quantum analogue of an instanton [25] possessing the energy ϵ .

The first equation in this system is the Schrödinger equation for the YM Hamiltonian

$$\hat{H} = \int d^3x \frac{1}{2}[E^2 + B^2]; \quad E = \frac{\delta}{i\delta A}.$$

The second one expresses the normalization assumed in the instanton YM model [25] at which the electric field E is transverse [11]:

$$\nabla_i E^i \Psi_\epsilon[A] = 0.$$

At last, the third equation implicates the *raising operator* [1]

$$T_1|n\rangle = |n+1\rangle$$

(with the winding number 1 as its eigenvalue). In the terminology [9], the transverse electric field E remains invariant with respect to all the “large” transformations T_1 , while the instanton wavefunction $\Psi_\epsilon[A]$ is manifestly covariant with respect to these transformations.

The raising operator T_1 can be represented explicitly in the shape [20]

$$T_1 = \exp\left(\frac{d}{dX[A]}\right) = \exp\left\{\left[\int d^3x B^2 \frac{g^2}{16\pi^2}\right]^{-1} \int d^3x B_i^a \frac{\delta}{\delta A_i^a}\right\},$$

with $X[A]$ being the YM winding number functional (its look is well known and we will not cite it here).

The above system of equations gives a correct definition (cf. [31]) of the θ -vacuum as a pseudomomentum operator possessing the common system of eigenfunctions $\{\Psi_\epsilon[A]\}$ with the momentum operator $\nabla_i^{ab}(A)$.

Indeed, one encounters the following problem with this definition [20, 31] of the θ -vacuum. The thing is that the operators $\hat{H}$ and T_1 don't commute (as it was argued in [20]) unless $\epsilon = 0$, i.e. the have no common eigenfunction $\Psi_\epsilon[A]$ at $\epsilon \neq 0$ (on the contrary, the Hamilton operator $\hat{H}$ commutes with the momentum $\nabla_i^{ab}(A)$).

It is obvious [20, 29, 30] that at setting $\epsilon = 0$, the definition [20, 31] of the θ -vacuum remains valid for imaginary as well as for real values of θ . This allows to represent θ as a complex number [30] $\theta = \theta_1 + i\theta_2$. In particular, at purely imaginary values of $\theta = \theta_2$ and $\epsilon = 0$, there exists the plane wave

$$\Psi_0[A] = \exp\{i(2\pi k + \theta_2)X[A]\} \equiv \exp\{iP_{\mathcal{N}}X[A]\} \quad (k \in \mathbf{Z})$$

which satisfies formally our definition [20, 31] of the θ -vacuum at

$$P_{\mathcal{N}} = 2\pi k + \theta_2 = 2\pi k \pm i8\pi^2/g^2.$$

Here the real part of $P_{\mathcal{N}}$, θ_1 , runs formally over the discrete set $2\pi k$, while its imaginary part $\theta_2 = 8\pi^2/g^2$ is continuous.

But this plane wave functional diverges manifestly at the $-$ sign before $8i\pi^2/g^2$ (this fact was pointed out, for instance, in the papers [4, 30]).

The name “quasimomentum” for θ has the following origin. The thing is that even real values of the topological momentum $P_{\mathcal{N}}$ have rather a fictive nature. This is so since the θ -term in the instanton YM effective Lagrangian (3.5) does not alter [8] the YM equations of motions

$$D_{\mu}F^{\mu\nu} = 0.$$

It is obvious that the first degree of the quasimomentum θ in the instanton YM effective Lagrangian (3.5)

implies its manifest P (and CP because of this) covariance.

The question about the “separate” C-covariance of the θ -item in (3.5) is more delicate. To understand which a “delicacy” is concealed here let us now resort to the arguments [26].

In this paper the effect influence the (real) θ -angle upon the dyon charge was investigated.

To determine the concrete effect CP violation of the dyon charge by θ -angle (latter proves to be conserved if $\theta = 0$; we have made sure in this above), one must apply a semi-classical analysis. For instance, the author [26] utilize the simple semi-classical analysis has been performed in the work [32].

In this work a semi-classical quantization of classical dyonic solutions. In a gauge in which fields disappear at the (spatial) infinity, classical dyonic solutions are periodic in time. The semi-classical quantization condition comes to the claim [26, 32] that $S + ET$ (the action during the time period T plus the energy times the time) should be an integer multiple of 2π .

Let I being the action per unit time. Then the above claim “that $S + ET$ should be an integer multiple of 2π ” is reduced to the relation

$$T(I + E) = 2\pi n \quad (n \in \mathbf{Z}).$$

The classical period T and the “abbreviated action” [26] $I + E$ (at $T = 1$) were calculated in Ref. [32] in the absence of the CP violation. It was found that

$$I + E = cq^2; \tag{3.6}$$

$$T = \frac{2\pi}{ec} \frac{1}{q}, \tag{3.7}$$

where q is the charge of the dyon and c is a constant¹⁰.

Simultaneously, one can construct (repeating the arguments [20, 29]; see also [30]) the family of purely real solutions for the topological momentum $P_{\mathcal{N}}$ satisfying the Schrödinger equation at $\epsilon = 0$ being parallel (via the θ) the eigenvalue of the raising operator T_1 :

$$P_{\mathcal{N}}^R = 2\pi k \pm 8\pi^2/g^2.$$

Issuing from this Eq., one can equate $\theta_1 = \pm 8\pi^2/g^2$ and assume θ_1 to vary in the interval $[-\pi, \pi]$; it is just that *real* θ -angle considered in modern gauge physics (see, for instance, [1]).

On the other hand, the presence of imaginary (i.e. space-like) momentum modes $8i\pi^2/g^2$ in the $P_{\mathcal{N}}$ spectrum (with the appropriate “diverged” plane waves) gives it impossible to give the correct probability description of the (topologically degenerated) θ -vacuum since the Hilbert space $\Psi_0^{(n)}[A]$ of its states becomes non-separable in this case.

The remarkable property of the real part $P_{\mathcal{N}}$ spectrum is following [20, 30]. It is obvious that such real topological momentum $P_{\mathcal{N}}^R$ vanishes (i.e. the instanton YM configuration stops) in the limit $g \rightarrow \infty$ for the YM coupling constant g . It is just the infrared QCD confinement limit as it is understood in modern physics. In the terminology [20] this case is referred to as the *infrared catastrophe*.

Note that purely imaginary (and thus space-like and unphysical) values $P_{\mathcal{N}} = \pm 8\pi^2 i/g^2$ (at $k = 0$) have no relation to the infrared catastrophe.

The results we have demonstrated now can be treated [20] as the presence of unphysical solutions to the Schrödinger equation at the application of ordinary quantization methods to a topologically nontrivial theory. In the paper [11] this statement was referred to as the so-called *no-go theorem*.

¹⁰It is not a large problem to calculate this constant, but a simple arguments shows that the same constant c appears in Eqs. (3.6) and (3.7).

The condition $T(I + E) = 2\pi n$ now gives

$$q = ne, \quad (3.8)$$

so that dyons possess integer charges as one expects in the absence of the CP violation.

Let us assume now that $\theta \neq 0$ and let us repeat the above calculations. At $\theta \neq 0$ the equations of motion are unchanged [8, 30], and there no change in the period T or the energy E . However, there is an extra contribution to the action I from the extra item $\Delta\mathcal{L}$ in the effective instanton YM Lagrangian (3.5)¹¹:

$$I + E = cq^2 + ceq \frac{\theta}{2\pi}. \quad (3.9)$$

The semi-classical (Bohr-Sommerfeld) quantization of $T(I + E)$ then gives

$$q = ne - \theta e / 2\pi. \quad (3.10)$$

So the allowed values of magnetic monopoles (if latter ones are contemplated in the model: for instance if Higgs modes are present in such model, violating the initial $SU(2)$ gauge symmetry) and are not integer if $\theta \neq 0$. In particular, there does not exist an electrically neutral magnetic monopole in that case.

If no magnetic and electric charges are presented in the gauge theory (i.e. there are no magnetic monopoles in that theory): for instance in the “pure” YM theory without Higgs modes (the instanton model is just such a case), one would set $q = 0$ in (3.10) (this does not contradict the Dirac quantization condition (2.1) since one deals with the uncertainty $0 \cdot 0$ in that case). Then

$$n = [\theta/2\pi], \quad (3.11)$$

i.e. the integer part of the number $\theta/2\pi$. It is, in fact, the ordinary connection between the θ -angle and the integer topological number n in the instanton model.

Now we should recall that in the absence of magnetic monopoles the θ -angle results no “true” motions in the YM theories. Instead of this, the θ -dependence in a YM model comes to purely tunnelling effects connected with instantons [1, 25] (for the topological number $n = 1$ such effects of the order $\exp(1/\alpha)$; $\alpha \equiv g^2/4\pi$).

¹¹In the light of our above conjecture [20, 29, 30] that $\theta = \theta_1 + i\theta_2$, the expression below, recast to the look

$$E = cq^2 - I + ceq \frac{\theta}{2\pi} = cq^2 - I + ceq \frac{\theta_1 + i\theta_2}{2\pi},$$

shows transparently that the “complex” θ -vacuum is not stationary. Due to the ordinary quantum mechanic canons, the life time of a θ -vacuum state with the energy E (indeed, we should set $E = 0$ since only such θ -vacuum states are compatible with the definition, as given above, of this vacuum) is

$$\tau = \frac{2\pi\hbar}{\theta_2}$$

(appropriately, its line width is $\theta_2/2\pi$).

This means that the θ -vacuum state with the energy $E = 0$ decays into (two) states: say, 1 and 2, for which $E_1 + E_2 = 0$. But these θ -vacuum states involving nonzero energies E_1 and E_2 are badly specified. As we have emphasized above, repeating the arguments [20], these values of energy cannot be simultaneously common eigenvalues of the YM Hamilton operator $\hat{H}$ and the raising operator T_1 .

In this is the next in turn contradiction about the YM instanton model. And moreover, the life time τ of a θ -vacuum state with the energy $E = 0$ is a finite number as $\theta_2 \neq 0$. Thus the $E = 0 \rightarrow E_1 + E_2$ vacuum decay occurs in the finite time τ . This implies once again bad specifying the quantum states $|1\rangle$ and $|2\rangle$ corresponding to the energies E_1 and E_2 as those not referring to the time infinities.

As it was discussed in Ref. [30] (repeating the said in the monograph [33]), in that case $\tau \neq \pm\infty$ it is impossible to describe correctly Feynman diagrams referring to the above $E = 0 \rightarrow E_1 + E_2$ vacuum decay since only at the claim $\tau \rightarrow \pm\infty$ the interaction representation of the system of quantum fields, set with the aid of the appropriate scattering matrix S , is true. If $\tau \neq \pm\infty$, one cannot pick out (as a consequence of the Haag theorem [33]) a correct Fock representation for interacting (quantum) fields.

But if magnetic monopoles (for instance, of the 't Hooft-Polyakov type [22, 23]) are incorporated in the YM theory, in the monopole sector of that theory there are classically allowed motions, dyons, with nonzero

$$n \sim \int d^4x F_{\mu\nu} \tilde{F}^{\mu\nu} \in \mathbf{Z}.$$

As a result, the θ -dependence in the monopole sector is something another than that connected with instantons: in particular, it is of the leading order rather different from $\exp(1/\alpha)$.

What happens, asks the author [26], in the YM theories in which CP is violated by another mechanism than θ ?

The fact that at $\theta = 0$ the dyons have integer charges is associated with the fact that $I + E$ in Eq. (3.6) is quadric in q , with no linear term. A linear term as in Eq. (3.9) leads to noninteger charges: the “non-integrality” is directly proportional to the coefficient of the linear term.

CP forbids such linear term since q is odd under CP. If CP is violated, regardless the violation mechanism, a linear term can be present.

Even if a linear term is absent on the classical level (for instance, if only couplings to fermions violate CP, since fermions do not enter classical solutions), it can be present on the quantum level due to loop corrections. Roughly speaking, one should recalculate $I + E$ from the *quantum* effective action (rather from the *classical* action). If CP is violated, loop corrections to the effective action would induce a term linear in q in the effective $I + E$ and therefore to cause monopole charges (if exist) to be not quite integer.

The same effect CP violating occurs, as we have demonstrated in the previous section, in the YMH model involving BPS magnetic monopoles (without any θ -dependence). The crucial point here is the presence of the BPS ansatz f_1^{BPS} , (1.8), in that model (as it can be seen from (2.16)). But these BPS magnetic monopole solutions induce (as it can be demonstrated) rather tree than loop Feynman diagrams.

4 Discussion.

The Dirac fundamental quantization [34] of the YMH model involving vacuum BPS monopole solutions (coming, as we have explained above, to solving the Gauss law constraint in terms of topological Dirac variables) implies some refining us said in the previous section about the CP violation in the presence of those vacuum BPS monopole solutions.

As it was discussed in the recent paper [30], resolving the YM Gauss law constraint

$$\frac{\delta W}{\delta A_0^a} = 0 \iff [D^2(A)]^{ac} A_{0c} = D_i^{ac}(A) \partial_0 A_c^i. \quad (4.1)$$

in terms of topological Dirac variables

$$\hat{A}_i^{(n)}(t, \mathbf{x}) = \hat{\Phi}_i^{(n)}(\mathbf{x}) + \hat{A}_i^{(n)}(t, \mathbf{x}); \quad n \in \mathbf{Z}; \quad \hat{A}_\mu = g \frac{A_\mu^a \tau_a}{2i\hbar c} \quad (4.2)$$

(here $\hat{A}_i^{(n)}(t, \mathbf{x})$ are perturbation excitations, *multipoles*, over the BPS monopole vacuum); satisfying the Coulomb gauge $D^i \hat{A}_i^{a(n)} = 0$, results the homogeneous equation

$$[D^2(A)]^{ac} \hat{A}_{0c} = 0 \quad (4.3)$$

for temporal components $\hat{A}_{0c}$ of YM fields.

As it was shown in Ref. [20], the solutions to the “homogeneous” YM Gauss law constraint (4.3) can be found in the shape

$$A_0^c(t, \mathbf{x}) = \dot{N}(t) \Phi_{(0)}^c(\mathbf{x}) \equiv Z^c, \quad (4.4)$$

implicating the topological dynamical variable $\dot{N}(t)$ and Higgs (topologically trivial) vacuum Higgs BPS monopole modes $\Phi_0^a(\mathbf{x})$.

Issuing from *zero mode solutions* [20] Z^a to the YM Gauss law constraint (4.1), it is easy to write down F_{i0}^a components of the YM tension tensor, taking the shape of so-called vacuum "electric" monopoles [3, 4]

$$F_{i0}^a = \dot{N}(t) D_i^{ac}(\Phi_k^{(0)}) \Phi_{0c}(\mathbf{x}).$$

In turn, vacuum "electric" monopoles F_{i0}^a generate the action functional

$$W_N = \int d^4x \frac{1}{2} (F_{0i}^c)^2 = \int dt \frac{\dot{N}^2 I}{2} \quad (4.5)$$

involving the *rotary momentum* [12]

$$I = \int_V d^3x (D_i^{ac}(\Phi_k^0) \Phi_{0c})^2 = \frac{4\pi^2 \epsilon}{\alpha_s} = \frac{4\pi^2}{\alpha_s^2} \frac{1}{V \langle B^2 \rangle}. \quad (4.6)$$

This action functional (describing the solid rotations of the YMH vacuum suffered the Dirac fundamental quantization) is manifestly Poincare (P) invariant since it can be recast to the look

$$W_N = \int dt \frac{P_N^2(t)}{2I}; \quad P_N = \dot{N}I. \quad (4.7)$$

Herewith

$$P_N = \dot{N}I = 2\pi k + \theta; \quad \theta \in [-\pi, \pi]; \quad (4.8)$$

is the momentum corresponding to the abovementioned solid rotations of the YMH vacuum. As it can be seen from (4.8), the P_N spectrum is purely real and thus rotary trajectories for the YMH vacuum suffered the Dirac fundamental quantization belong to the physical ones.

It is just the momentum spectrum free from imaginary, i.e. tachionic, modes, unlike the instanton case, as discussed in Section 3.

Besides that the action functional (4.7) is manifestly P-invariant, it is also C-invariant. To understand this fact, we should compare Eq. (4.7) (it is a definite functional of the YMH BPS monopole vacuum rotary energy $P_N^2(t)/2I$) with Eq. (3.9) [26].

In the latter case the appropriate energy $I + E$ associated with the instanton θ vacuum proves to be linear (i.e. odd) by e and θ . This just implies CP violating.

The former case is rather different. As it follows from (4.6), (4.7), the action functional (4.7) is directly proportional to the vacuum "electric" field ("electric" monopole) F_{i0}^a , induced, in turn, by a Higgs vacuum BPS monopole mode $\Phi_{0c}(\mathbf{x})$.

As it was shown in Ref. [12] (see also [30]),

$$F_{i0}^a \equiv E_i^a = \dot{N}(t) (D_i(\Phi_k^{(0)}) \Phi_{(0)})^a = P_N \frac{\alpha_s}{4\pi^2 \epsilon} B_i^a(\Phi_{(0)}) = (2\pi k + \theta) \frac{\alpha_s}{4\pi^2 \epsilon} B_i^a(\Phi_{(0)}). \quad (4.9)$$

Latter relations were got with the aid of the Bogomol'nyi equation (1.6). Since the YM coupling constant $\alpha_s \equiv g^2/4\pi(\hbar c)^2$ enters Eq. (4.9), it is already squared by g , while the rotary momentum I , (4.6), is of the order g^4 .

But at accepting the normalization (2.12) [8] for the electric charges q (referring to the Higgs Bose condensate), the vacuum "electric" field F_{i0}^a will be of the order q^{-2} (respectively, the YMH BPS monopole vacuum rotary energy $P_N^2(t)/2I$ will be of the order q^{-4}). Thus the YMH model with vacuum BPS monopoles quantized by Dirac is C-invariant as that possessing the action functional (4.7) even by the total electric charge $Q = \sum_i q_i$ of (topologically degenerated) Higgs vacuum BPS monopole modes $\Phi_{c(n)}(\mathbf{x})$ [11].

This us demonstrated CP conservation is the next in turn remarkable feature of that model.

References

- [1] T. P. Cheng, L.- F. Li, Gauge Theory of Elementary Particle Physics, 3rd edn. (Oxford University Press 1988).
- [2] L. H. Ryder, Quantum Field Theory, 1st edn. (Cambridge University Press 1984).
- [3] L. D. Lantsman, V. N. Pervushin, The Higgs Field as The Cheshire Cat and his Yang-Mills "Smiles", Proc. of 6th International Baldin Seminar on High Energy Physics Problems (ISHEPP), Dubna, Russia, 10-15 June 2002; [arXiv:hep-th/0205252];
L. D. Lantsman, Minkowskian Yang-Mills Vacuum, [arXiv:math-ph/0411080].
- [4] L. D. Lantsman, V. N. Pervushin, Yad. Fiz. **66**, 1416 (2003) [Physics of Atomic Nuclei **66**, 1384 (2003)]; [arXiv:hep-th/0407195].
- [5] L. D. Lantsman, Superfluid Properties of BPS Monopoles, [arXiv:hep-th/0605074].
- [6] M. K. Prasad, C. M. Sommerfeld, Phys. Rev. Lett. **35**, 760 (1975);
E. B. Bogomol'nyi, Yad. Fiz. **24**, 449 (1976).
- [7] R. Akhoury, J.-H. Jun, A. S. Golghaber, Phys. Rev. **D 21**, 454 (1980).
- [8] A. S. Schwarz, Kvantovaja Teorija Polja i Topologija, 1st edition (Nauka, Moscow, 1989) [A. S. Schwartz, Quantum Field Theory and Topology (Springer, 1993)].
- [9] L. D. Faddeev, in Proc. of 4th Int. Symp. on Nonlocal Quantum Field Theory, Dubna, USSR, 1976, JINR D1-9768, p. 267;
R. Jackiw, Rev. Mod. Phys. **49** (1977) 681.
- [10] V. N. Pervushin, Dirac Variables in Gauge Theories, Lecture Notes in DAAD Summerschool on Dense Matter in Particle and Astrophysics, JINR, Dubna, Russia, August 20- 31, 2001; Fiz. Elem. Chast. Atom. Yadra **34**, 679 (2003) [Phys. Part. Nucl. **34**, 348 (2003)]; [hep-th/0109218].
- [11] A. M. Khvedelidze, V. N. Pervushin, Helv. Phys. Acta, **67**, 637 (1994).
- [12] D. Blaschke, V. N. Pervushin, G. Röpke, Topological Gauge invariant Variables in QCD, MPG-VT-UR 191/99, [arXiv:hep-th/9909193].
- [13] D. Blaschke, V. N. Pervushin, G. Röpke, Topological Invariant Variables in QCD, in Proceeding of the Int. Seminar Physical variables in Gauge Theories, Dubna, September 21-25, 1999, edited by A. M. Khvedelidze, M. Lavelle, D. McMullan and V. Pervushin (E2-2000-172, Dubna, 2000), p. 49, [arXiv:hep-th/0006249].
- [14] T. T. Wu, C. N. Yang, Phys. Rev. **D 12**, 3845 (1975).
- [15] S. G. Matinyan, G. K. Savidy, Nucl. Phys. **B 134**, 539 (1978); A. A. Vladimirov, D. V. Shirkov, Usp. Fiz. Nauk **129**, 407 (1979) [Sov. Phys. Usp. **129**, 860 (1979)].
- [16] D. J. Gross, F. Wilczek, Phys. Rev. Lett. **30**, 1343 (1973); Phys. Rev. **D 8**, 3633 (1973);
H. D. Politzer, Phys. Rev. **D 8**, 3636 (1973).
- [17] N. N. Bogoliubov, J. Phys. **9**, 23 (1947);
N. N. Bogoliubov, V. V. Tolmachev, D. V. Shirkov, Novij Metod v Teorii Sverchprovodimosti, 1st edition (Izd-vo AN SSSR, Moscow, 1958, pp. 5-9).
L. D. Landau, JETP **11**, 592 (1941); DAN USSR **61**, 253 (1948).
- [18] I. M. Khalatnikov, Teorija Sverxtekychesti, 1st edition (Nauka, Moscow, 1971).

- [19] L. D. Landau, E. M. Lifschitz, Lehrbuch der Theoretischen Physik (Statistische Physik, Band 5, teil 2), in German, 1st edition by H. Escrig and P. Ziesche (Akademie-Verlag, Berlin, 1980).
- [20] V. N. Pervushin, Teor. Mat. Fiz. **45**, 394 (1980) [Theor. Math. Phys. **45**, 1100 (1981)].
- [21] V. G. Levich, Yu. A. Vdovin, V. A. Mjamlin, Kurs Teoreticheskoy Fiziki, v. 2, 2nd edition (Nauka, Moscow, 1971).
- [22] G. 't Hooft, Nucl. Phys. **B 79**, 276 (1974).
- [23] A. M. Polyakov, Pisma JETP **20**, 430 (1974) [Sov. Phys. JETP Lett. **20**, 194 (1974)]; Sov. Phys. JETP Lett. **41**, 988 (1975).
- [24] F. Bruckmann, G. 't Hooft, Phys. Rep. **142**, 357 (1986), [arXiv:hep-th/0010225].
- [25] A. A. Belavin, et al., Phys. Lett. **59**, 85 (1975);
R. Jackiw , C. Rebbi, Phys. Lett. **B 63**, 172 (1976);
C. G. Jr. Callan, R. F. Dashen, D. J. Gross, Phys. Rev. **D 17**, 2717 (1977).
- [26] E. Witten, Phys. Lett. **B 86**, 283 (1979).
- [27] P. A. M. Dirac, Proc. Roy. Soc. **A 133**, 69 (1931).
- [28] D. Zwanziger, Phys. Rev. **176**, 1480 (1978);
J. Schwinger, Phys. Rev. **144**, 1087 (1968); Phys. Rev. **173**, 1536 (1968).
- [29] A. S. Galperin, V. N. Pervushin, JINR P2-11830 (1978).
- [30] L. D. Lantsman, Nontrivial Topological Dynamics in Minkowskian Higgs Model Quantized by Dirac., [arXiv:hep-th/0610217].
- [31] C. G. Jr. Callan et al., Phys. Rev. **D 17**, 2717 (1978).
- [32] E. Tomboulis, G. Woo, Nucl. Phys. **B 107**, 221 (1976).
- [33] N. N. Bogoliubov, A. A. Logunov, A. I. Oksak, I. T. Todorov, Obshie Prinzipi Kvantovoj Teorii Polja, 1st edition (Nauka, Moscow, 1987).
- [34] P. A. M. Dirac, Proc. Roy. Soc. A 114 (1927) 243; Can. J. Phys. 33 (1955) 650.